

Stars of WIMPs

F. Pisano* and J. L. Tomazelli†

Instituto de Física Teórica

Universidade Estadual Paulista

Rua Pamplona, 145

01405-000-São Paulo, SP

Brazil

Abstract

A class of boson-fermion stars, whose spin-0 and spin-1/2 constituents interact through a U(1) current-current term in the Lagrangian density, is analysed. It is shown that it describes the low-energy behavior of a system of weakly interacting massive particles (WIMPs) from the leptonic sector of the minimal supersymmetric standard model. In this case the effective coupling constant λ is related to the Fermi constant G_F .

PACS numbers: 12.60 – Models beyond the standard model, 93.35 – Dark matter

Typeset using REVTeX

*Supported by FAPESP

†Supported by CAPES

Along the last two decades there has been an increasing interplay between gravity and elementary particle theories, in which the former has found a field-theoretical support such as in the inflationary models [1]. These models predict that the density parameter of the Universe $\Omega_{\text{total}} = 1$, in contrast with $\Omega_B \lesssim 0.15$ for the baryonic density of the standard Big Bang nucleosynthesis. This indicates the existence of nonbaryonic dark matter. Furthermore, there is a direct evidence from astronomical observations of groups of galaxies and clusters, whose dynamics indicate the presence of hidden matter [2]. Among the candidates for the dark matter are boson stars [3] and composite structures of bosons and fermions [4], which might be formed in the early Universe. However, none of the models considered up to now succeeded in taking into account any specific interaction among the constituents of such stellar objects. Recently it was proposed a current-current type interaction between those bosons and fermions [5]. Here we reinterpret this interaction in the context of the minimal supersymmetric standard model (MSSM) as an effective interaction corresponding to the exchange of a massive gauge boson and determine the strength of the coupling. In this case the fermion and scalar particles are massive charged leptons together with their respective supersymmetric partners, a particular set of weakly interacting massive particles (WIMPs) which are the prime candidates to the above mentioned nonbaryonic dark matter.

Let us start with the Lagrangian density for pure gravity and the free complex scalar field without self-coupling,

$$\mathcal{L} = \frac{R}{16\pi G} + \partial_\mu \Phi^* \partial^\mu \Phi + m_s^2 \Phi^* \Phi , \quad (1)$$

where

$$\Phi(r, \tau) = \phi(r) e^{-i\omega\tau} \quad (2)$$

and ϕ is a complex scalar field.

The background fermion field is described by a perfect fluid [6] with energy density ρ and pressure p :

$$\rho = K(\sinh t - t) ,$$

(3)

$$p = \frac{K}{3} \left(\sinh t - 8 \sinh \frac{t}{2} + 3t \right) ,$$

where t is a parameter, $K = m_f^4/32\pi^2$, and m_f is the fermion mass.

The corresponding energy-momentum tensor for the free fields reads as

$$T_{\mu\nu}^{(0)} = T_{\mu\nu}^s + T_{\mu\nu}^f , \quad (4)$$

with

$$T_{\mu\nu}^s = \partial_\nu \Phi^* \partial_\mu \Phi + \partial_\mu \Phi^* \partial_\nu \Phi - g_{\mu\nu} (\partial_\lambda \Phi^* \partial^\lambda \Phi + m_s^2 \Phi^* \Phi) , \quad (5)$$

$$T_{\mu\nu}^f = (\rho + p) u_\mu u_\nu - p g_{\mu\nu} , \quad (6)$$

where indices s and f label the scalar and fermion fields, respectively.

The complete Lagrangian density includes the contact interaction term

$$\mathcal{L}^{\text{int}} = -\lambda j_\mu(\psi) J^\mu(\Phi) , \quad (7)$$

where

$$j_\mu(\psi) = \bar{\psi} \gamma_\mu \psi , \quad (8)$$

$$J^\mu(\Phi) = i(\Phi^* \partial^\mu \Phi - \Phi \partial^\mu \Phi^*) . \quad (9)$$

It is invariant under global U(1) gauge transformations and contributes to the energy-momentum tensor with

$$T_{\mu\nu}^{\text{int}} = -\lambda (J_\mu j_\nu - g_{\mu\nu} j_\alpha J^\alpha) . \quad (10)$$

In the region of low frequencies the fermion-current can be replaced by

$$j_\mu(\psi) = \bar{\psi} \Gamma_\mu \psi , \quad (11)$$

where

$$\Gamma_0 \equiv u_0 \quad , \quad \Gamma_i \equiv i u_i . \quad (12)$$

The four-vector $u_\mu = (u_0, u_r, u_\theta, u_\varphi)$ is the four-velocity of the fermion fluid in spherical coordinates. Our metric convention is $(+, -, -, -)$ otherwise specified.

Let us pick out from the interaction Lagrangian density for the broken supersymmetric $SU(2) \otimes U(1)$ theory [7] those terms that contribute to the scattering of massive leptons and scalar-leptons, which can be identified with the effective interaction Lagrangian density (7) in the low-energy limit:

$$\mathcal{L}_{\tilde{l} \rightarrow l \tilde{l}}^{\text{int}} = \mathcal{L}_{llV} + \mathcal{L}_{\tilde{l}\tilde{l}V} , \quad (13)$$

with

$$\mathcal{L}_{llV} = -\frac{g}{2 \cos \theta_W} \bar{l} \gamma_\mu (g_V - g_A \gamma_5) l Z^\mu + e \bar{l} \gamma_\mu l A^\mu , \quad (14)$$

$$\mathcal{L}_{\tilde{l}\tilde{l}V} = -i \frac{g}{2 \cos \theta_W} \tilde{l}^{*T} \overleftrightarrow{\partial}_\mu (\tilde{g}_V - \tilde{g}_A \Gamma_5) \tilde{l} Z^\mu + ie \tilde{l}^{*T} \overleftrightarrow{\partial}_\mu \tilde{l} A^\mu , \quad (15)$$

where we have defined

$$\Gamma_5 \equiv \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} , \quad \tilde{l} \equiv \begin{pmatrix} \tilde{l}_L \\ \tilde{l}_R \end{pmatrix} \quad (16)$$

and

$$g_V = \tilde{g}_V \equiv -\frac{1}{2} + 2 \sin^2 \theta_W \quad \text{and} \quad g_A = \tilde{g}_A \equiv -\frac{1}{2} . \quad (17)$$

These terms give rise to the interaction between a fermion- and a scalar-current mediated by a vector gauge boson. Since (7) is a contact interaction, the exchanged virtual particle must be massive. Thus, neither the last term in (14) nor that in (15) must be taken into account, since they produce a long-range interaction. The scalar neutral-currents that comprise the interaction Lagrangian density (15) were written in a suggestive fashion in order to stress the correspondence with the respective fermion counterparts in (14), where $\tilde{l}_L$ and $\tilde{l}_R$ stand for the supersymmetric partners of the ordinary charged lepton states $l = \{e, \mu, \tau\}$ projected in the chiral basis.

For the moment, we consider processes involving electrons and scalar-electrons. In the MSSM such scalar particles must correspond to fermions in a definite chirality state. The amplitudes for the $e^- \tilde{e}_R^- \rightarrow e^- \tilde{e}_R^-$ and $e^- \tilde{e}_L^- \rightarrow e^- \tilde{e}_L^-$ processes in momentum space are

$$\mathcal{M}^R = -ig^2 \frac{\tan^2 \theta_W}{2} \bar{u}(\vec{p}_1') (\not{p}_2 + \not{p}_2') (g_V - g_A \gamma_5) u(\vec{p}_1) \frac{1}{k^2 - M_Z^2}, \quad (18)$$

$$\mathcal{M}^L = -ig^2 \frac{1 + 2 \sin^2 \theta_W}{4 \cos^2 \theta_W} \bar{u}(\vec{p}_1') (\not{p}_2 + \not{p}_2') (g_V - g_A \gamma_5) u(\vec{p}_1) \frac{1}{k^2 - M_Z^2}, \quad (19)$$

where $k^2 = (p_1' - p_1)^2$ and we have written the massive gauge boson propagator in the Feynman gauge.

At the infrared, where spin effects become negligible, we can safely disregard the piece of (18) and (19) which contains the axial-current. In fact, it follows from the Dirac equation

$$(\not{p} - m)u_r(\vec{p}) = 0; \quad r = 1, 2$$

for positive-energy four-spinors and relation

$$\sigma^{ij} = -\gamma^0 \gamma^5 \gamma^k; \quad k = 1, 2, 3 \quad \text{in cyclic order}$$

that

$$\bar{u}_s(\vec{p}') \gamma^\mu \gamma^5 u_r(\vec{p}) = \bar{u}_s(\vec{p}') \gamma^\mu \left[\frac{p_0 - m \gamma^0}{|\vec{p}|} \right] (-1)^{r+1} u_r(\vec{p}), \quad (20)$$

where $u_r(\vec{p})$ are eigenstates of the helicity operator

$$\sigma_{\vec{p}} = \frac{\vec{\sigma} \cdot \vec{p}}{|\vec{p}|}.$$

Making the replacement $\gamma^\mu \rightarrow \Gamma^\mu$, where the Γ 's were defined in (12), we immediately see that the axial-current vanishes.

Hereafter we shall consider only the vector-current amplitudes, which for $k^2 \ll M_Z^2$ reduce to

$$\mathcal{M}^R = i \frac{g^2}{4M_Z^2} \tan^2 \theta_W (-1 + 4 \sin^2 \theta_W) \bar{u}(\vec{p}_1') (\not{p}_2 + \not{p}_2') u(\vec{p}_1), \quad (21)$$

$$\mathcal{M}^L = i \frac{g^2}{4M_Z^2} \frac{(1 + 2 \sin^2 \theta_W)(-1 + 4 \sin^2 \theta_W)}{2 \cos^2 \theta_W} \bar{u}(\vec{p}_1') (\not{p}_2 + \not{p}_2') u(\vec{p}_1). \quad (22)$$

The ratio between the total cross-sections constructed from amplitudes (21) and (22) is given by

$$\frac{\sigma_R}{\sigma_L} = \left(\frac{2x}{1+2x} \right)^2, \quad (23)$$

with $x \equiv \sin^2 \theta_W$. Since the value of x decreases with energy, the first process is suppressed in the infrared limit, where we have

$$\frac{\sigma_R}{\sigma_L} \sim 4x^2 - 16x^3 + 48x^4 - \dots, \quad x < 1.$$

If we compare the amplitude which stems from the effective interaction (7) with (22), we extract the value

$$\begin{aligned} \lambda &= \frac{g^2}{4M_Z^2} \frac{(1 + 2\sin^2 \theta_W)(1 - 4\sin^2 \theta_W)}{2\cos^2 \theta_W} \\ &= \frac{G_F}{4\sqrt{2}} (1 + 2\sin^2 \theta_W)(1 - 4\sin^2 \theta_W), \end{aligned} \quad (24)$$

where G_F is the Fermi constant.

The effective coupling constant λ can be expressed in terms of the adimensional coupling $\bar{\alpha}$ that appears in the Einstein equations for the metric coefficients and the Klein-Gordon equation for the radial component of the scalar field [5]

$$\lambda = \frac{m_s \bar{\alpha}}{2\bar{n}_f}, \quad (25)$$

where $\bar{n}_f$ is the average fermion density. From (24) and (25) we have

$$\bar{\alpha} = \frac{G_F}{2\sqrt{2}} \frac{\bar{n}_f}{m_s} (1 + 2\sin^2 \theta_W)(1 - 4\sin^2 \theta_W). \quad (26)$$

All the scalar-leptons have essentially the same lower bound for their masses, $m_s \sim 45$ GeV [8], so that $\bar{\alpha}$ is independent of the scalar-lepton flavor. For a background of a degenerate electron gas [9], $\bar{n}_f$ is of order 10^{30} cm^{-3} . Substituting the values for G_F , x and m_s , we obtain

$$\bar{\alpha} \lesssim 10^{-19} \quad (27)$$

and therefore we do not reach the critical value for the second-order phase transition reported in [5], since the electron density is not high enough to create bound states of electrons and scalar-electrons.

Summing up we have investigated gravity in the presence of scalar fields in interaction with a background fermion fluid, in the framework of a fundamental particle theory, namely the MSSM in which the Bose-Fermi system can be interpreted as a self-gravitating system of WIMPs, the strongest candidates to the dark matter. This enabled us to attribute a precise meaning to the arbitrary coupling constant λ of the effective theory, which results to be proportional to the Fermi constant G_F and the electroweak mixing parameter $\sin^2 \theta_W$. For a degenerate Fermi gas the value of the adimensional coupling $\overline{\alpha}$ is below the critical value for Bose-Einstein condensation.

REFERENCES

- [1] A. H. Guth, Phys. Rev. **D 23**, 347 (1981); A. D. Linde, Phys. Lett. **B 108**, 347 (1982);
A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
- [2] G. C. Stewart, C. R. Canizares, A. C. Fabian and P. E. J. Nilsen, Astrophys. J. **278**, 53
(1984).
- [3] R. Ruffini and S. Bonazzola, Phys. Rev. **187**, 1767 (1969).
- [4] A. B. Henriques, A. R. Liddle and R. G. Moorhouse, Phys. Lett. **B 233**, 99 (1989).
- [5] C. M. G. de Sousa and J. L. Tomazelli, preprint gr-qc/9507043.
- [6] S. Chandrasekhar, Astrophys. J. **140**, 417 (1964).
- [7] H. E. Haber and G. L. Kane, Phys. Rep. **C 117**, 75 (1985).
- [8] L. Montanet *et al.* (Particle Data Group), Phys. Rev. **D 50**, 1173 (1994).
- [9] Ya. B. Zeldovich and I. D. Novikov, *Relativistic Astrophysics*, Vol.1, The University of
Chicago Press (1971).